

Heavy metal-induced oxidative stress on seed germination and seedling development: a critical review

Mihiri Seneviratne · Nishanta Rajakaruna · Muhammad Rizwan ·
H. M. S. P. Madawala · Yong Sik Ok · Meththika Vithanage

Received: 8 November 2016 / Accepted: 26 June 2017
© Springer Science+Business Media B.V. 2017

Abstract Heavy metal contamination in soils can influence plants and animals, often leading to toxicosis. Heavy metals can impact various biochemical processes in plants, including enzyme and antioxidant production, protein mobilization and photosynthesis. Hydrolyzing enzymes play a major role in seed germination. Enzymes such as acid phosphatases, proteases and α -amylases are known to facilitate both seed germination and seedling growth via mobilizing nutrients in the endosperm. In the presence of heavy metals, starch is immobilized and nutrient sources become limited. Moreover, a reduction in proteolytic enzyme activity and an increase in protein and amino

acid content can be observed under heavy metal stress. Proline, is an amino acid which is essential for cellular metabolism. Numerous studies have shown an increase in proline content under oxidative stress in higher plants. Furthermore, heat shock protein production has also been observed under heavy metal stress. The chloroplast small heat shock proteins (Hsp) reduce photosynthesis damage, rather than repair or help to recover from heavy metal-induced damage. Heavy metals are destructive substances for photosynthesis. They are involved in destabilizing enzymes, oxidizing photosystem II (PS II) and disrupting the electron transport chain and mineral metabolism.

M. Seneviratne
Department of Botany, Faculty of Natural Sciences, Open
University of Sri Lanka, Nawala, Nugegoda, Sri Lanka

N. Rajakaruna
Unit for Environmental Sciences and Management,
North-West University, Potchefstroom, South Africa

N. Rajakaruna
Biological Sciences Department, California Polytechnic
State University, San Luis Obispo, CA 93407, USA

M. Rizwan
Department of Environmental Sciences and Engineering,
Government College University, Allama Iqbal Road,
Faisalabad 38000, Pakistan

H. M. S. P. Madawala
Department of Botany, University of Peradeniya,
Peradeniya, Sri Lanka

Y. S. Ok (✉)
Korea Biochar Research Center & School of Natural
Resources and Environmental Science, Kangwon National
University, Chuncheon 24341, Korea
e-mail: soilok@kangwon.ac.kr

M. Vithanage (✉)
Environmental Chemodynamics Project, National
Institute of Fundamental Studies, Kandy, Sri Lanka
e-mail: meththikavithanage@gmail.com

M. Vithanage
Office of the Dean, Faculty of Applied Sciences,
University of Sri Jayewardenepura, Nugegoda, Sri Lanka

Although the physiological effects of Cd have been investigated thoroughly, other metals such as As, Cr, Hg, Cu and Pb have received relatively little attention. Among agricultural plants, rice has been studied extensively; additional studies are needed to characterize toxicities of different heavy metals on other crops. This review summarizes the current state of our understanding of the effects of heavy metal stress on seed germination and seedling development and highlights informational gaps and areas for future research.

Keywords Antioxidant system · Starch mobilization · Proline · Protein degradation · Chlorophyll · Heavy metal stress

Introduction

Chemical waste resulting from industrial activities and agricultural effluents such as fertilizer, herbicides and pesticide has contributed to the increased accumulation of heavy metals in soil (Fu et al. 2014; Hu et al. 2013; Nicholson et al. 2003; Neilson and Rajakaruna 2015). Since heavy metals are non-degradable, they can cause long-term deleterious effects on ecosystem health. Copper (Cu), As and Cd are the major heavy metals present in agrochemicals while Hg, Cr, Pb and As are rich in industrially contaminated soils (Wuana and Okieimen 2011). Cadmium (Cd), Hg, Pb, Cu and As are some of the most toxic heavy metals or metalloids found in soil (Dago et al. 2014; Clemens 2006; Hadi et al. 2013). Cadmium, for example, is nonessential for plant metabolic activities and a known phytotoxicant (Gratão et al. 2005). The long biological half-life of Cd contributes to its mutagenic, cytotoxic and carcinogenic properties (Sánchez-Virosta et al. 2015; Waalkes 2003). Ore smelting, coal burning, industrial effluents, and As-containing herbicides and insecticides release As, a toxic metalloid, to the environment (Lin et al. 2016). Mercury (Hg), which is one of the most potent toxic heavy metals, is released mainly from mining activities (Lacerda 1997; De Lacerda and Salomons 2012); however, low amounts can also be found in agrochemicals (Saeueia et al. 2013; Falkowska et al. 2013). Copper is released from several industrial activities and by the

excess usage of fungicides. Copper-containing fungicides and bactericides are used extensively for disease control in numerous crops. Lead (Pb), a metal that can remain in the soil for thousands of years (Kumar et al. 1995), can be released to the environment via mining activities (Bakirdere et al. 2016) and other industrial processes (Bénard et al. 2014). Although some heavy metals are considered as micronutrients (Cu, Zn, Mn, etc.), other metals and metalloids are highly toxic (Hg, Pb, Cd, As, etc.) and play no role in plant nutrition and their presence in high concentrations is highly toxic (Nagajyoti et al. 2010; Boyd and Rajakaruna 2013).

The organic matter and clay content and pH are the major factors governing the availability of heavy metals in soils (Zeng et al. 2011; Spurgeon and Hopkin 1996). A high proportion of humified organic matter (OM), i.e., mature compost, can also decrease the bioavailability of heavy metals, while increasing the residual fraction of heavy metals in the soils (Castaldi et al. 2005), which is mainly attributed to the precipitation caused by the complex formation (Bolan and Duraisamy 2003). Moreover, the large specific surface area of clay minerals contributes to higher immobilization of heavy metals in soil (Prost and Yaron 2001; Usman et al. 2005). As the soil pH is increased, the immobilization is also increased for many heavy metals, especially for those metals present as cations (i.e., Pb, Cd and Cr). Biochar application to soil, which increases the soil pH, has also demonstrated the effect of pH in heavy metal immobilization (Herath et al. 2015).

High concentrations of heavy metals in soil cause a number of deleterious effects on plants (Khan et al. 2000), such as growth retardation, destruction of chlorophyll, disorders in biochemical activities, mutations and reproductive disorders (Schützendübel and Polle 2002; Sharma and Agrawal 2005; Gall and Rajakaruna 2013). Seed germination, one of the most significant stages in a plant's life, is sensitive to chemical and physical conditions of the rhizosphere (Bewley 1997). Although the seed coat can act as a principle barrier limiting harmful effects of heavy metals, most seeds and seedlings show a decline in germination and vigor in response to heavy metal stress (Adrees et al. 2015), causing a major concern for agricultural and forestry practices. Therefore, the effects of heavy metals on germination and seedling growth are an important research area deserving

extensive study. Recent studies have documented that via inhibition of storage food mobilization, reduction in radical formation, disruption of cellular osmoregulation and the degradation of proteolytic activities, heavy metals cause inhibition of seed germination and seedling development (Adrees et al. 2015; Barceló and Poschenrieder 1990; Perfus-Barbeoch et al. 2002; Karmous et al. 2015; Baszyński 2014) (Fig. 1). However, several management practices have been utilized to minimize heavy metal stress on plants grown in contaminated soils. While physical and chemical technologies are useful in remediation of heavy metal contaminated soils (Dermont et al. 2008; Wu et al. 2010), the use of different amendments is one of the low-cost and efficient methods utilized to minimize the plant heavy metal uptake in contaminated soils; biochar, crab shells and organic matter are a few such amendments (Uchimiya et al. 2010). Amendments are used to immobilize the contaminants, thereby limiting the bioavailable fraction (Bandara et al. 2016, 2017; Herath et al. 2017; Kumarathilaka and Vithanage 2017). In addition to amendments, there are also internal mechanisms in place, which can limit the interaction of heavy metals in plant metabolism. Phytochelatins and metallothioneins are such proteins that may play vital roles in heavy metal detoxicity in microorganisms and plants (Shen et al. 2010).

This review focuses on the influence of heavy metals and metalloids (especially, Cd, Hg, Ag and As) on seed germination and seedling development in plants, focusing particularly on physiological and biochemical effects on germinating seeds and developing seedlings.

Effect of heavy metals on seed germination and seedling growth

Heavy metals are known to negatively influence seed germination and seedling development (Table 1). Cadmium leads to germination inhibition as documented by several studies. For example, when *Sorghum bicolor* (Poaceae) seeds were treated with a Cd concentration series (0.0, 0.5, 1.0, 2.0 and 3.0 mM), a significant reduction in germination was observed in the presence of the metal (Barceló and Poschenrieder 1990). The decrease in germination appears to result from the inhibition of physiological and metabolic activities of the seed. Since Cd is able to decrease the water stress tolerance of plants (Barcelo et al. 1986), it can cause a loss of turgor pressure at a higher relative water content compared to that of non-treated plants. The effect of Cd on plant water status has been examined with germinating rice seeds (Barceló and Poschenrieder 1990), showing that the water status of germinating rice seeds is highly




Fig. 1 Different effects of heavy metals on seed germination and seedling development

Table 1 A summary of studies examining the effects of heavy metals on seed germination and seedling development

Heavy metal	Species	Findings	References
Al	<i>Zea mays</i>	Reduction in seed germination was observed beyond 50 mg L ⁻¹	Nasr (2013)
As	<i>Brassica oleracea</i> , <i>Amaranthus</i> sp., <i>Raphanus sativus</i> , <i>Daucus carota</i>	Reduction in germination was observed at 10 mg L ⁻¹	Dutta et al. (2014)
Cd	<i>Triticum aestivum</i>	Cd showed toxicity at 5 mg L ⁻¹ in root and shoot growth. Seed germination and germination energy were affected at 20 mg L ⁻¹ which is aggravated by further addition of Cd from 50 to 80 mg L ⁻¹	Ahmad et al. (2012)
	<i>Oryza sativa</i>	Cd shows a significant reduction in rice seed germination index, vigor index, root and shoot lengths as well as fresh weight compared to control at 100 µM	He et al. (2014)
	<i>Picea omorika</i>	1 mM concentration inhibited germination	Prodanovic et al. (2016)
	<i>Solanum nigrum</i>	Germination rate was dramatically reduced at 200–300 µmol L ⁻¹ Cd	Liu et al. (2012)
	<i>Triticum aestivum</i>	A decrease in germination was recorded from 20 mg Cd L ⁻¹	Ahmad et al. (2013)
	<i>Helianthus annuus</i>	Gradual reduction in germination (2–10 mg L ⁻¹)	Imran et al. (2013)
	<i>Triticum aestivum</i>	Seed germination was stimulated at low concentrations (0–2.5 mg L ⁻¹) while a reduction in germination was observed under high concentrations (5–30 mg L ⁻¹)	Mahdieh et al. (2013)
	<i>Vigna radiata</i> and <i>Glycine max</i>	1 mg kg ⁻¹ As addition stimulated seed germination and increased about 12% of the germination weight, but the seed germination was significantly suppressed when with As addition was over 5 mg kg ⁻¹	Wan et al. (2013)
	<i>Festuca rubra</i>	No germination inhibition was observed at 25 mg L ⁻¹ while a reduction in germination was observed at 25 and 50 mg L ⁻¹	Vázquez de Aldana et al. (2014)
	<i>Brassica juncea</i>	No significant effect was observed in seed germination up to 250 µM	Srivastava et al. (2013)
Cd, Pb, Cu	<i>Arachishypogeeae</i>	Concentrations of 75 and 100 mg L ⁻¹ , Cd and Pb, respectively, decreased seed germination while Cu condensed seed germination at 100 mg L ⁻¹	Abraham et al. (2013)
Cr	<i>Cucumis melo</i>	Reduction in seed germination as the concentration increased	Akinci and Akinci (2010)
Co, Cr	<i>Phaseolus vulgaris</i>	Reduction in seed germination at high concentrations (10 ⁻² M)	(Zeid 2001)

Table 1 continued

Heavy metal	Species	Findings	References
Cu	<i>Oryza sativa</i>	The germination of seeds decreased as the concentration increased (0.2–1.5 mM)	Ahsan et al. (2007a)
	<i>Triticum aestivum</i>	The germination of seeds decreased from 5 mg L ⁻¹	Singh et al. (2007)
	<i>Zea mays</i>	Increased seed germination at 0.1 mM	Bashmakov et al. (2005)
	<i>Triticum aestivum</i> and <i>Oryza sativa</i>	Reduced germination by more than 35 and 60%, respectively, with 10 μM	Mahmood et al. (2007)
	<i>Medicago sativa</i>	Inhibit seed germination by 39% with 40 mg L ⁻¹	Aydinalp and Marinova (2009)
	<i>Eruca sativa</i>	0.3–1.2 mM did not decrease seed germination while lower Cu concentration (<0.7 mM) increased seed germination	Zhi et al. (2015)
Hg	<i>Vignaradiata</i>	No significant reduction in seed germination was shown at 1 mM; however, beyond 3 mM, a reduction in germination was observed	Muhammad et al. (2015)
	<i>Albizia lebbek</i>	A significant gradual reduction in seed germination was observed beyond 1 mM	Iqbal et al. (2014)
	<i>Cajanus cajan</i>	Reduction in seed germination was observed beyond 5 mg L ⁻¹	Patnaik and Mohanty (2013)
	<i>Brassica napus</i>	Reduction in seed germination was observed beyond 1 mM	Rezaei et al. (2013)
Ni	<i>Medicago sativa</i>	40 ppm inhibited seed germination by 24%	Aydinalp and Marinova (2009)
	<i>Zea mays</i>	Reduction in seed germination was observed beyond 50 mg L ⁻¹	Nasr (2013)
Pb	<i>Triticum aestivum</i>	Increased germination rate and root mass with Pb concentration less than 50 mg L ⁻¹	Kang et al. (2009)
	<i>Lens culinaris</i>	A significant reduction in germination was observed from 0.5 mM	Cokkizgin and Cokkizgin (2015)

affected by Cd. Since water imbibition is a major requirement for seed germination, the negative influence of Cd on the water content of the seed is significant (Barceló and Poschenrieder 1990). Similarly, an inhibition of root and coleoptile growth has been observed in seedlings under Cd stress (Kuriakose and Prasad 2008). Cadmium appears to cause an inhibition of carbohydrate hydrolysis and the translocation of hydrolyzed sugars, resulting in a reduction in seedling growth. Mercury (Hg) is also known to negatively influence seed germination and seedling growth (Muhammad et al. 2015). As *Vigna radiata* (Fabaceae) was treated with a Hg concentration series (1, 3, 5 and 7 mM), there was no significant reduction

under 1 mM; however, the 7-mM treatment showed the highest reduction among all treatments, resulting in 42, 70, 66 and 47% reduction in seed germination, seedling length, root length and seedling dry weight, respectively. Though silver (Ag) has often been used in nanofertilizers, only a handful of studies have been conducted to assess the effects of Ag with respect to seedling growth. A study examining the effects of polyvinylpyrrolidone-coated silver nanoparticles (PVP-AgNP) on 11 wetland plants (*Lolium multiflorum*, *Panicum virgatum*, *Carex lurida*, *C. scoparia*, *C. vulpinoidea*, *C. crinita*, *Eupatorium fistulosum*, *Phytolacca americana*, *Scirpus cyperinus*, *Lobelia cardinalis* and *Juncus effusus*) showed that Ag

nanoparticles had no effect on germination, while Ag 40 mg L⁻¹ gum arabic coated silver nanoparticle (GA-AgNP) exposure significantly reduces the germination rate in three species (*S. cyperinus*, *J. effusus* and *P. americanum*) and enhances the germination rate of one (*E. fistulosum*) species (Yin et al. 2012). Arsenic (As), another toxic metalloid, is known to suppress plant growth (Han et al. 2015), showing significant negative effects on root and shoot length and root and shoot biomass of wheat seedlings; the growth of root and shoot was inhibited by high concentration of As (5–20 mg kg⁻¹), where the inhibition of root was stronger than that of the shoot (Wang et al. 2002). A reduction in starch granules in seeds reflects the mobilization of insoluble sugars; scanning electronic microscopic images can be used to analyze the immobilized starch content in seedlings (Singh et al. 2003); Kuriakose and Prasad (2008) have observed that under 0.5 mM Cd, starch granules further decrease, indicating more mobilization of starch. Whereas most of the starch were in seeds treated with 2.0 mM Cd, a significant reduction of acid phosphates was observed in both seed and seedling with 2.0 and 3.0 mM of Cd, respectively. Activity staining using densitometric scanning showed the reduction of an isozyme in all Cd-treated seedlings, whereas a reduction in total protease activity was observed in both embryonic axes and seeds compared to the control. Further, a significant decrease in rice seed germination index, vigor index, root and shoot lengths as well as fresh weight compared to control was observed in plants exposed to Cd (He et al. 2014). The study showed that the exogenous nitric oxide donor sodium nitroprusside (SNP) can balance the inhibitory effects of rice seed germination and seedling growth.

Changes in antioxidant system

Cadmium is a toxic metal influencing seedling growth and development. A concentration series of Cd was used to determine Cd stress on the antioxidant system of rice seedlings (Ali et al. 2002). The fluctuation of antioxidant concentrations was different at varying Cd levels. Super oxide dismutase (SOD) and glutathione reductase (GR) showed an increasing pattern in both leaves and roots. Ascorbate peroxidase (APX) and peroxidase (POD), however, showed an increase at low concentrations and a decline at high

concentrations. Further, catalase (CAT) activity showed contrasting effects, with low concentrations in leaves and high concentrations in roots. A similar pattern of non-enzymatic and enzymatic antioxidants was detected with oilseed rape treated with Cd (Yan et al. 2015). Interestingly, interactive effects of Cd and Pb in rice seedlings cause a reduction in glutathione and ascorbate activity and their redox ratios, while causing an increase in activities of antioxidative enzymes, superoxide dismutase (SOD) and guaiacol peroxidase (Srivastava et al. 2014).

Acetyl salicylic acid (ASA) is another protective measure utilized by plants for removing oxygen free radicals. Under low As concentrations, the ASA content increases, whereas with an increase in As concentrations, the amount of oxygen free radicals increases (Li et al. 2007). With further increase in As concentration, the ASA content decreases in wheat seedling leaves in order to eliminate the oxygen free radicals (Li et al. 2007). Ascorbate peroxidase (APX) activity decreased slightly at low concentrations of As (0–5 mg kg⁻¹) and then increased as the concentration of As increased. The CAT activity displayed an increasing trend at lower concentrations (0–1 mg kg⁻¹) and a decreasing trend with increasing As concentrations (5–20 mg kg⁻¹). However, POD activity increased significantly with an increase in As concentrations. The enhanced production of antioxidant enzymes represents typical defense mechanism against As accumulation and toxicity (Li et al. 2007). Although many studies have shown changes in antioxidant systems with respect to heavy metals, the mechanisms of heavy metal influence on the antioxidants have not been explored. Additional studies should be carried out for a better understanding of antioxidant production under oxidative stress. Table 2 summarizes studies examining the effects of heavy metals on antioxidant properties of germinating seed.

Production of plant growth regulators

Plant growth regulators play an important role in seed germination. Gibberellic acid (GA) is a plant hormone controlling seed germination (Groot and Karssen 1987). Abscisic acid (ABA) reduces this effect while cytokinins are able to reverse the effect of ABA (AA Khan 1968). Heavy metals can influence endogenous plant growth regulator (PGRs) levels in seed

Table 2 Summary of studies showing heavy metal-induced antioxidant production in germinating seeds and seedlings

Metal	Species	Changes in antioxidant system	References
As	<i>Triticum aestivum</i>	The ascorbate peroxidase and glutathione <i>S</i> -transferase activities increased with the increase in As concentration, while glutathione reductase (GR) activities increased only at 0.25 mM. The monodehydroascorbate reductase and catalase activities were not changed upon exposure to As. The activities of dehydroascorbate reductase and glyoxalase I decreased at any level of As, while glutathione peroxidase and glyoxalase II activities decreased only upon 0.5 mM As	Hasanuzzaman and Fujita (2013)
As–Se	<i>Vigna radiata</i>	As severely inhibited the growth at 10 μM. The As-treated (10 μM) plants supplemented with 5 μM Se showed improved growth indicating an antagonistic interaction between the two elements. This likely resulted from elevated levels of enzymatic superoxide dismutase, catalase, ascorbate peroxidase and glutathione reductase	Malik et al. (2012)
As	<i>Oryza sativa</i>	Superoxide dismutase, catalase, ascorbate peroxidase and glutathione reductase increased in arsenite treated seedlings, while dehydroascorbate reductase activity declined initially during 5–10 days and then increased	Mishra et al. (2011)
As	<i>Oryza sativa</i>	Up-regulation of superoxide dismutase, ascorbate peroxidase and glutathione reductase with exposure to As	Shri et al. (2009), Rai et al. (2011)
	<i>Oryza sativa</i>	Up-regulation of superoxide dismutase, ascorbate peroxidase and glutathione reductase with exposure to As	Shri et al. (2009), Rai et al. (2011)
Ag	<i>Ricinus communis</i>	The peroxidase activity and superoxide dismutase activity increased with increasing Ag concentration	Yasur and Rani (2013)
Ag–Cd	<i>Vicia faba</i>	In leaves, catalase activity declined, but peroxidase activity increased with time. In roots, catalase activity and peroxidase activity increased during early stages and then declined. Superoxide dismutase activity showed an upward trend with an increase in the duration of treatment after 3 and 6 days and then declined both in leaves and in roots (at 9 days) under 1–50 μM	Zhang et al. (2009)
Cd	<i>Picea omorika</i>	1 mM inhibited the activities of catalase, superoxide dismutase and peroxidase	Prodanovic et al. (2016)
	<i>Trigonella foenum-graecum</i>	With concentrations greater than 0.1 mM, catalase, superoxide dismutase and ascorbate peroxidase were increased in different parts of the plant	Zayneb et al. (2015)
	<i>Brassica napus</i>	From 100 μm catalase, superoxide dismutase, glutathione reductase and ascorbate peroxidase are reduced	Mohamed et al. (2012)
	<i>Oryza sativa</i>	Marked elevation in superoxide dismutase and peroxidase activities was observed with increase in Cd. The activity of catalase increased in seedlings grown at moderately toxic Cd (100 μM) level, whereas a highly toxic Cd (500 μM) level inhibited catalase activity.	Shah et al. (2001)
	<i>Brassica juncea</i>	Peroxidase, catalase and superoxide dismutase in the leaves of plants increased significantly in parallel with an increase in the concentration of the metal	Irfan et al. (2014)
Cd, Pb	<i>Oryza sativa</i>	Superoxide dismutase and guaiacol peroxidase increased in metal-treated seedlings. Catalase activity increased during early stages (8 days) of metal exposure and declined by 16 days	Srivastava et al. (2014)
Cr	<i>Glycine max</i>	The shoots contain higher catalase activity than roots. The reduction of catalase was observed from 5 mg L ⁻¹ onwards, and it was severely affected at 200 mg L ⁻¹ . The peroxidase and super oxide dismutase activity increased with the increase in Cr	Sankarganesh et al. (2015)
Cr	<i>Brassica juncea</i> and <i>Vigna radiata</i>	Higher activities of superoxide dismutase, ascorbate peroxidase, catalase and glutathione reductase were observed in shoot of <i>B. juncea</i> and <i>V. radiata</i> at all levels of Cr treatments (0, 50, 100, 200 μM)	Diwan et al. (2010)
Cu, Zn	<i>Cumumis sativus</i>	Significant increase in superoxide dismutase, peroxidase and catalase activities at 100 mg L ⁻¹ metal concentrations	Kim et al. (2012)

Table 2 continued

Metal	Species	Changes in antioxidant system	References
Cu	<i>Helianthus annuus</i>	In the presence of 0.5 mM metal concentration, superoxide dismutase activity was increased while catalase, ascorbate peroxidase, glutathionine reductase were decreased	Gallego et al. (1996)
Mn	<i>Oryza sativa</i>	3 and 6 mM Mn increased superoxide dismutase guaiacol peroxidase and catalase in 10 day old seedlings, but declined by 20 days under Mn treatment. Moreover, ascorbate peroxidase monodehydroascorbate reductase, dehydroascorbate reductase and glutathione reductase increased significantly with Mn	Srivastava and Dubey (2011)
Pb, Cd and Hg	<i>Lycopersicon esculentum</i>	50 μ M decreased catalase activity in roots. The activity of ascorbate peroxidase was unchanged in roots and in stems, while it was diminished in leaves. However, guaiacol peroxidase activity was increased only in roots and stems	Mazhoudi et al. (1997)
Ni and Zn	<i>Cajanus cajan</i>	Increased the superoxide dismutase, peroxidase and glutathione reductase and reduced catalase activity in all Zn and Ni treatments	Rao and Sresty (2000)

germination. The mevalonic acid pathway is the main biosynthetic pathway for GA production (Arteca 1996). Several studies report an increase in GA content during seed germination (Debeaujon and Koornneef 2000; Nambara et al. 1991). The GA content showed a reduction in chickpea seeds in the presence of Zn and Pb (1.0–10 mM) and Cd. Although GA is reduced under high Zn concentrations, it showed an increase at low Zn (0.1 mM) concentrations (Atici et al. 2005). The mechanism behind heavy metal-caused reduction in GA production is also not well understood.

Abscisic acid is another PGR governing the seed germination process. It is largely responsible for seed dormancy and germination (Fujii et al. 2007). The role of ABA on inhibition of seed germination been well studied (Koornneef et al. 1984). It is known that ABA levels in plants are enhanced with exposure to environmental stress (Jiang and Zhang 2002). An increase in ABA concentration was observed in Cd-tolerant *Oryza sativa* (Poaceae) seedlings while an exogenous application of ABA increased Cd tolerance in *O. sativa* seedlings (Hsu and Kao 2003). Further, ABA causes a reduction in transpiration rates, helping to reduce shoot Cd accumulation, thereby reducing heavy metal-caused oxidative stress (Salt et al. 1995). Similarly, both GA and ABA are able to reduce translocation of heavy metals in *O. sativa* although an external application of these hormones was unable to overcome oxidative stress (Rubio et al. 1994). However, ABA appears to mediate paclobutrazol-induced Cd tolerance in *O. sativa* (Hsu and Kao 2005).

Effect of heavy metals on α -amylase activity

Hydrolyzing enzymes which are produced by the aleurone layer of seed play a major role in seed germination (Yan et al. 2014; Taiz and Honigman 1976). Enzymes such as acid phosphatases (ACPs), proteases and α -amylases facilitate both seed germination and successful seedling establishment and growth via mobilizing food reserves in the endosperm (Yan et al. 2014). In the presence of Cd and Cu, α -amylase, acid phosphatase and alkaline phosphatase activities were reduced in the endosperm of barley seeds (Kalai et al. 2014). Similarly, as germinating barley seeds were soaked under various concentrations of arsenite and arsenate, a reduction of α -amylase was detected (Sanal et al. 2014).

Starch, proteins and fat are the reserved food sources in a seed, with starch as the most abundant food source of most angiosperm seed (Dua and Sawhney 1991). Starch hydrolyzation can occur either by means of an amylolytic or phosphorolytic pathway. Amylases are the major enzymes responsible for the starch mobilization process. As pea (*Pisum sativum*; Fabaceae) seeds were treated with a Cd concentration series, both α - and β amylase activities were suppressed (Chugh and Sawhney 1996). After 7 days of germination, the total amylase activity with 0.25, 0.50 and 1.00 mM of Cd was 56, 42 and 33%, respectively. This suppression was more pronounced in the α -amylase activity compared to the β -amylase activity. However, the net effect was due to the synergistic effect of both enzymes, with α -amylase degrading

starch into smaller oligosaccharides, creating non-reducing ends for exo-hydroxylation of β -amylase (Fulton et al. 2008). Therefore, the starch degradation can be impeded even at low Cd levels due to the reduction in α -amylase activity.

Lead is another toxic heavy metal released by various industrial activities. A 50% reduction of α -amylase was observed in pea seedlings treated with Pb, resulting in high starch content. The decline in 6-phosphogluconate dehydrogenase (G6PDH) production and an increase in hexokinase (HXK) production in roots and shoots of stressed seedlings signify that hexoses could be utilized for glycolysis more than the pentose phosphate pathway (Devi et al. 2013). Similarly, starch immobilization by Pb via reduction of starch hydrolyzing enzymes has been shown in *Brassica campestris* (Singh et al. 2011).

Chromium also reduces seed germination by causing deleterious effects on hydrolyzing enzymes (Dua and Sawhney 1991). With the increase in Cr concentrations, the total amylase activity showed a reduction in germination pea seeds; however, on the 8th day since exposure, the level of activity was comparable to that of the control (Dua and Sawhney 1991). Therefore, it was hypothesized that the toxic effect can be mitigated with a prolonged exposure to the metal. Similarly, α -amylase showed a more pronounced reduction in concentrations than β -amylase with an increase of Cr, suggesting that the recovery activity of β -amylase plays an important role in the restoration of total enzyme activity. Furthermore, Cr has inhibitory effects on acid phosphatases phytase and ribonuclease, with no significant effects on amylase and protease (Dua and Sawhney 1991).

Protein catabolism

Proteins are another source of energy for seed germination. Proteins in seeds are found as both storage substances and enzymes. Germinating seeds growing under heavy metals stress show alterations and behavior of many enzymes. Extensive studies have been conducted on the effect of As, Cd, Al and Ni on enzyme activities (Maheshwari and Dubey 2008, 2009; Mishra and Dubey 2006; Mishra et al. 2011; Shah and Dubey 1995, 1998; Shah et al. 2001). During seed development and maturation, proteins are synthesized and placed within membrane-bound protein bodies. During seed germination, the storage proteins are hydrolyzed

and transported to the growing points of the embryo. Proteases and peptides hydrolyse proteins into a mixture of free amino acids and small peptides (Ranki et al. 1990). The profile of a cell can be altered by external stress signals. Proteolytic enzyme changes with respect to Cd stress have been studied in rice seeds (Shah and Dubey 1998). Cadmium concentration were increased in the rice seeds with a reduction in protease activity that were subjected to heavy metals and at the same time, enhanced uptake of Cd in embryo axes than in the endosperm (Shah and Dubey 1998). Additionally, there was a 30–50% decline in leucine amino peptidase activity, which catalyzes the hydrolysis of leucine residues, and a marked increase in carboxypeptidase activity, which catalyzes the release of arginine or lysine residues from polypeptides in endosperm. However, there was always a reduction in peptidase activity in embryo axes of germinating seeds (Shah and Dubey 1998).

Karmous et al. (2012, 2014a) examined the effects of Cu on protease activity in seeds. The effects of Cu on bean (*Phaseolus vulgaris*) cotyledons were assessed with respect to protease activity (Karmous et al. 2012). The enzymes involved in storage protein mobilization during bean germination have also been studied via electrophoresis. Copper influenced both embryo development and total protease activity. Moreover, cysteine, aspartic and metalloprotease activities were all decreased under Cu stress, with an increase in the serine-protease activity. Hence, excess Cu appears to disturb the release of nitrogen from storage tissue during seed germination. It has been suggested that there may be a relationship between defense capacity and proteolytic responses in Cu-treated embryos (Karmous et al. 2014b).

Similar results were obtained with rice seedlings treated with As (Mishra and Dubey 2006). RNA and protein degradation are key processes in cellular biochemistry. RNase is one of the key marker enzymes for stress conditions. The changes in the activity of RNase are an indicator of stress conditions (Shah and Dubey 1995). Since rice is sensitive to As, the effects of As₂O₃ on activities of RNase, protease, peptidases as well as on the levels of RNA, proteins, free amino acids and proline in growing rice seedlings were tested. Compared to the control, RNA proteins and proline were increased. A reduction in hydrolysis of RNA and proline in rice seeds could be due to As induced inhibition of RNase and protease activity. Moreover,

protease and ribonuclease inhibition of rice seedling has been documented under Ni stress (Maheshwari and Dubey 2008). There is acellular level balance of macromolecules such as RNA and proteins within their synthesis and degradation. A possible suppression of RNA hydrolysis in rice plants under Cd toxicity has also been documented (Shah and Dubey 1995). This may be due to the reduced activity of RNase under metal stress, hindering the degradation of RNA. Since the cellular RNA levels and protein synthesis are main processes that maintain cellular development, metal stress can disturb the cellular biochemistry (Shah and Dubey 1995).

Abiotic stresses can induce or suppress different forms of proteases. A relationship between the enzyme activities of protein metabolism and the level of free amino acids/proteins was reported in As-treated plants (Mishra and Dubey 2006). There was a suppression of both protease and leucine aminopeptidase activity while an enhanced activity was observed in carboxypeptidase in rice under As stress, showing the variation in the response of different enzymes under heavy metal stress. Moreover, carboxypeptidase activity showed a considerable tolerance under the concentration range of As used.

Protein damage is one of the crucial effects of oxidative stress. Oxidation of residual amino acid side chains and formation of covalent cross-linkages between proteins can denature cellular proteins. The rapid elimination of those proteins is necessity to avoid toxicities during cell metabolism. The ubiquitin (Ub) proteasome system is an important non-lysosomal proteolytic pathway involved in the degradation of intracellular proteins under biotic and abiotic stress (Davies 2001; Jackson and Durocher 2013). An increase in protein carbonylation and an decrease in the reduced S–H pool reflect protein damage in bean seeds under heavy metal stress (Kranner and Colville 2011). Moreover, Cu showed an induced inactivation of Ub-proteasome, which limits the removal of oxidatively damaged proteins in germinating bean seeds (Karmous et al. 2014a).

Protective role of proline against heavy metal stress

Proline is an amino acid which plays an important role in maintaining cellular homeostatis via osmotic regulation (Fariduddin et al. 2013; Iqbal et al. 2016).

Studies show that the proline content is increased under different environmental stresses. An accumulation of proline was observed under numerous biotic stresses (Fabro et al. 2004). It was hypothesized that proline is able to scavenge free radicals produced by a variety of abiotic stresses, including heavy metal stress. The potential of proline to scavenge the free radicals was assessed with in vitro assay systems, namely graft copolymerization and auto-oxidation of pyrogallol (Kaul et al. 2008).

Although the relationship between proline accumulation and abiotic stress tolerance in crops is always not obvious, studies have shown that there is a correlation between the plant proline accumulation and stress conditions (Ashraf and Foolad 2007). The accumulation of proline was tested in *Cajanus cajan*, *Vigna mungo* and *Triticum aestivum* under Cd, Co, Pb and Zn stress (Saradhi 1991). According to the phenotypic data obtained, Cd was the most toxic metal among the heavy metals used. An accumulation of proline was observed in all seedlings under heavy metal stress. The proline level was inversely proportional to the fresh weight of the plant, irrespective of the heavy metal, suggesting that materials used for plant growth were used for proline production. Cadmium appeared to be the most effective heavy metal in stimulating proline production (Saradhi 1991).

Free proline was also measured in response to Cu, Cd and Zn in non-tolerant and metal-tolerant *Silene vulgaris* (Schat et al. 1997). The constitutive proline concentration in the leaves of the metal-tolerant ecotype was 5–6 times greater than in the non-tolerant ecotype. Similarly, Ni-induced proline was also documented in rice (Maheshwari and Dubey 2009) and wheat (Gajewska and Skłodowska 2008). In addition to the metal, the plant species is also an important consideration in biochemical processes of plants. In *Silene vulgaris*, Cu was the most effective proline inducer (Schat et al. 1997). Complete inhibition of proline accumulation under heavy metal reduced transpiration explains the dependence of metal-induced proline accumulation on the development of metal-induced water deficit in leaves.

However, contradictory results have also been recorded on the water deficit hypothesis of proline. The accumulation of proline was tested in detached rice leaves treated with excess Cu (10 mM) (Chen et al. 2001). Both CuSO₄ and CuCl₂ were effective in simulating proline accumulation in rice leaves.

Interestingly, there was no effect on the relative water content of detached rice leaves with excess Cu, suggesting that there is no relationship between Cu-induced proline accumulation and water deficit condition in plants (Chen et al. 2001). Moreover, there was an increase in abscisic acid (ABA) concentration in detached rice leaves under Cu stress, which indicates the possibility of ABA to increase proline in stressed tissue. Similarly, an increment in proline content was observed with increasing Cu concentration in germinating wheat seedlings. There was also an increase in proline accumulation in the root of wheat seedlings even at a non-toxic concentration of Cu (1 μ M) (Chen et al. 2001).

Proline provides diverse protective functions in mitochondria. During cellular stress conditions, proline helps via supplying energy which is needed to recover from stress. It enhances reducing potential of mitochondria by donating electrons to the respiratory chain via oxidation by proline dehydrogenase (PDH) and proline-5-carboxylate dehydrogenase (P5CDH) proline (Szabados and Savouré 2010). The high proline accumulation under heavy metal stress has also been hypothesized as leading to a reduction in proline degradation, an increase in proline biosynthesis, a decrease in proline utilization in protein synthesis and the hydrolysis of proteins (Charest and Ton Phan 1990).

Heavy metal-induced protein changes

Proteins play an important role in all living systems. Any physiological or biochemical change, whether negative or positive, will be reflected by alterations in protein activity. Globulin is a major protein in plants, containing a high content of sulfur-containing amino acids (Kriz 1989; Xiang et al. 2013). Compared to that of untreated seeds, the storage protein accumulation was comparatively higher under heavy metal treatment. Seed proteome analysis reveals the presence of globulin and its precursor in cereals in that these proteins play an important role in basic cellular processes in rice embryogenesis and germination (Komatsu et al. 2005). The up-regulation of globulin-like proteins with their precursors was observed in the presence of excessive Cu in germinating rice (Ahsan et al. 2007a). Similarly, an up-regulation of the globulin precursors in germinating maize seeds was also detected with potassium dichromate, indicating

the activation of storage proteins (Labra et al. 2006). These results suggest that heavy metals may hinder the glycolytic pathways in germinating seeds, since the accumulation of seed storage proteins was higher relative to that of the untreated seeds.

The physiological, biochemical and proteomic changes in germinating rice seeds have also been tested in response to Cu stress. The total proteins were extracted and subjected to one-dimensional sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Differences were observed in the protein band patterns between the control and Cu-treated samples. The protein pattern changes were further analyzed by two-dimensional electrophoresis for better separation and characterization of germination-related proteins (Østergaard et al. 2004). The Cd-induced protein profile change was also examined in rice seeds with SDS-PAGE. Changes in protein patterns due to heavy metal binding proteins were visualized (Ahsan et al. 2007b), revealing results similar to those shown in previous studies with rice and maize (Ahsan et al. 2007a; Labra et al. 2006). Since the SDS-PAGE is not sensitive enough to determine the pin point changes of polypeptide patterns, the 2-DE analysis was used to investigate the changes in proteins (Zhang and Riechers 2004). A pre-fractionation technique was used to extract the rare proteins. The selected spots were subjected to MALDI-TOF mass spectrometry for peptide mass mapping, and 21 upregulated proteins were identified. These proteins were involved in RNA processing, signal transduction, protein biosynthesis, antioxidant defense and germination processes (Ahsan et al. 2007b).

Further, as *Lepidium sativum* plantlets were exposed to Cd, considerable amounts of unmodified or minimally modified storage proteins were detected in parallel to a reduction in plantlet elongation (Gianazza et al. 2007). This suggests the inhibition of storage protein catabolism with metal exposure during germination and seedling elongation. Since the *L. sativum* plantlets exhibit changes in protein expression with low concentrations of heavy metal, they are effective bio-indicators for assessing environment pollution.

Heat shock proteins in photosynthesis protection

Heat shock proteins are general stress proteins implicated in the protection of physiological processes

during most abiotic stresses (Basile et al. 2013). Chloroplast small Hsp (smHsp) is not a typical protein produced by plants in the absence of stress. Studies have confirmed the protective role of smHsps on photosynthetic electron transport (Ph_{et}), during heat, oxidative and photoinhibition stresses (Shakeel et al. 2012). As *Zea mays* (corn) plants were treated with varying soil concentrations of Cu, Ni, Pb and Zn, net photosynthesis was decreased with all heavy metals, with a greater decline shown under higher concentrations and during extended exposure times (Heckathorn et al. 2004). However, chloroplast small heat shock protein content was increased to levels comparable to levels induced by severe acute heat stress in the presence of Cu, Ni, Pb and Zn (Heckathorn et al. 2004). Further, the production of chloroplast small Hsp is a primary response in plants to heavy metal accumulation and its mode of protection is the limitation of photosynthesis damage, rather than repair or recovery from heavy metal damage (Downs et al. 1999; Waters et al. 1996). Hsp70 and small Hsp are induced in response to heavy metals, with Hsps protecting the cell membrane during heavy metal stress (Hall 2002).

Heavy metal-induced photosynthesis damage in seedlings

Heavy metals are often described as destructive substances for photosynthesis (Prasad and Strzałka 1999). They are involved in disrupting enzyme activities, oxidizing PS II and disorganizing the electron transport chain and mineral metabolism. A reduction in chlorophyll and heme levels was observed when germinating seedlings of *Phaseolus vulgaris* were treated with Cd (Somashekaraiiah et al. 1992). There was a significant increase in lipid peroxide levels in the presence of Cd. However, a dose-dependent induction of lipoxigenase activity was observed with the Cd^{2+} treatment. Moreover, a reduction in antioxidant enzyme activity was also seen with the Cd treatment. Inhibition of free radical scavenging enzymes via lipid peroxide accumulation may be a reason for the reduction in the chlorophyll and heme levels.

Similarly, a growth reduction, inhibition of chlorophyll content and photosynthesis have been reported in wheat in the presence of Cd (Ouzounidou et al. 1997). The indirect effect of Cd on essential nutrient

content in plants is one reason for photosynthesis inhibition. Moreover, structural damage of chloroplasts was evident by the distorted shape and the dilation of the thylakoid membranes. These ultrastructural changes suggest the possibility of Cd-induced premature senescence. However, Sheoran et al. (1990) demonstrated that the reduction in photosynthesis in the presence of Cd was due to the effect on stomatal conductance and not due to decreased enzyme activities. They show that Cd^{2+} inhibited the leaf O_2 evolution alongside a reduction in photosynthesis.

The chlorophyll molecule with Mg has the highest capacity of electron release from a single excited state (Küpper et al. 1996). As the central Mg atom of chlorophyll is substituted with heavy metals such as Hg, Cu, Cd, Ni, Pb, it prevents photosynthetic light-harvesting, contributing to a decline in photosynthesis (Mishra and Dubey 2005). Under low light irradiance, some of the chlorophyll substituted with heavy metals is more stable than Mg-containing chlorophyll. Consequently, crops can remain green even when they are dead. Even though several studies have been carried out to determine the effects of heavy metals on photosynthesis of plants, limited studies have been carried out on the effects of heavy metals on the photosynthesis of seedlings.

Reduction in heavy metal toxicity in seeds

The reduction in heavy metal toxicity in seeds can be indirectly achieved by immobilization of heavy metals in soil. This is mainly done with the use of different organic and inorganic amendments. Biochar, green waste manure, compost are some of the organic amendments that can be used to immobilize heavy metals (Branzini and Zubillaga 2012; Gadepalle et al. 2007; Park et al. 2011). Composted organic soil amendments have the ability to bind with heavy metals and can cause rapid immobilization. Due to the high surface area of carbonaceous sorbents such as black carbon and activated carbon with a high affinity for heavy metals, there can be an increased immobilization of heavy metals. Numerous studies have shown the efficiency of heavy metal adsorption of by biochar (Uchimiya et al. 2010; Inyang et al. 2012). Further, the addition of organic matter can also play a major role in heavy metal adsorption in soil (Davis 1984). Several studies have shown that the sorption power of organic matter is mainly due to its cation exchange capacity

(Basta et al. 1993). Therefore, the addition of these soil amendments can indirectly reduce the potential toxicity of heavy metals on seed.

Future perspectives

Heavy metal release to the environment is pronounced growing environmental problem, leading to detrimental effects at the ecosystem level. When early life cycle stages such as seed germination and seedling establishment are affected by heavy metals, there can be lasting consequences at the individual plant, population and community levels. The ways in which heavy metals can cause toxicities, including in storage food mobilization, photosynthesis and plant osmoregulation, can vary depending on the type of heavy metal, its speciation, soil pH, plant species and the mobility of the metal. In order to overcome heavy metal-induced toxicities, plants employ numerous mechanisms, including the activation of the antioxidant system, proline production, production of heat shock proteins and acetyl salicylic acid, among others. There are many informational gaps relating to the study of heavy metal influence on seed germination and seedling establishment. Some of these areas are summarized below to generate interest for further investigation.

- Protein hydrolysis and mobilization are key processes in seed germination. The interference of heavy metals with key germination-related proteins has not been well documented at the proteomic level. The study of proteomics under heavy metal stress can be a fruitful area for future study.
- The specific processes or cell components protected by Hsps remain largely unknown. Additionally, the specific Hsps induced by heavy metals and their cellular location also deserve further investigation.
- GA production by the embryo is a major step in seed germination. More studies are needed in the area of heavy metal influences on GA production.
- Ubiquitin (Ub) proteasome system is an important non-lysosomal proteolytic pathway involved in the degradation of intracellular proteins under biotic and abiotic stress. However, more research is necessary to document the involvement of the Ub

system in the degradation of specific cellular proteins. There are numerous studies carried out on the effect of heavy metals on hydrolyzing enzymes; however, there is a lack of studies on the effects of heavy metals on the seed's aleurone layer where the hydrolyzing enzymes are produced.

- Although it is known that heavy metals are able to reduce the water uptake in germinating seeds, the exact mechanisms for heavy metal-induced osmotic stress are unclear. Additional studies are needed to explore the mechanisms of heavy metal-induced osmoregulation in germinating seed.
- Most of the studies to date have been carried on cereal crops, especially on rice. Hence, it is imperative that additional studies are carried out on a broad range of plants, including fruit and vegetable crops and wild plants. Research on wild plant tolerance of heavy metals can also provide useful information on understanding heavy metal-induced stresses on germination and early seedling growth.
- The effects of understudied metals including Ag are important to examine in the laboratory as these metals are now commercially used as nano fertilizer.

Seed germination and early seedling growth are two important stages of plant development. An understanding of the multitude of ways in which heavy metals influence these critical stages in plant growth and development is fundamental for advancing plant science and developing better practices in agriculture, forestry and restoration. This review highlights the primary ways in which heavy metals impact critical stages of plant development in the hope of fostering additional research for improving our knowledge on the deleterious effects heavy metals have on seeds and seedlings and how such effects could be minimized with additional research in phytotechnologies.

References

- Abraham, K., Sridevi, R., Suresh, B., & Damodharam, T. (2013). Effect of heavy metals (Cd, Pb, Cu) on seed germination of *Arachis hypogaea* L. *Asian Journal of Plant Science and Research*, 3(1), 10–12.
- Adrees, M., Ali, S., Rizwan, M., Ibrahim, M., Abbas, F., Farid, M., et al. (2015). The effect of excess copper on growth and physiology of important food crops: A review.

- Environmental Science and Pollution Research*, 22(11), 8148–8162.
- Ahmad, I., Akhtar, M. J., Asghar, H. N., & Zahir, Z. A. (2013). Comparative efficacy of growth media in causing cadmium toxicity to wheat at seed germination stage. *International Journal of Agriculture & Biology*, 15, 517–522.
- Ahmad, I., Akhtar, M. J., Zahir, Z. A., & Jamil, A. (2012). Effect of cadmium on seed germination and seedling growth of four wheat (*Triticum aestivum* L.) cultivars. *Pakistan Journal of Botany*, 44(5), 1569–1574.
- Ahsan, N., Lee, D.-G., Lee, S.-H., Kang, K. Y., Lee, J. J., Kim, P. J., et al. (2007a). Excess copper induced physiological and proteomic changes in germinating rice seeds. *Chemosphere*, 67(6), 1182–1193.
- Ahsan, N., Lee, S.-H., Lee, D.-G., Lee, H., Lee, S. W., Bahk, J. D., et al. (2007b). Physiological and protein profiles alternation of germinating rice seedlings exposed to acute cadmium toxicity. *Comptes Rendus Biologies*, 330(10), 735–746.
- Akinci, I. E., & Akinci, S. (2010). Effect of chromium toxicity on germination and early seedling growth in melon (*Cucumis melo* L.). *African Journal of Biotechnology*, 9(29), 4589–4594.
- Ali, M. B., Chun, H. S., Kim, B. K., & Lee, C. B. (2002). Cadmium-induced changes in antioxidant enzyme activities in rice (*Oryza sativa* L. cv. Dongjin). *Journal of Plant Biology*, 45(3), 134–140.
- Arteca, R. N. (Ed.). (1996). Historical aspects and fundamental terms and concepts. In *Plant growth substances* (pp. 1–27). Berlin: Springer.
- Ashraf, M., & Foolad, M. (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environmental and Experimental Botany*, 59(2), 206–216.
- Atici, Ö., Açar, G., & Battal, P. (2005). Changes in phytohormone contents in chickpea seeds germinating under lead or zinc stress. *Biologia Plantarum*, 49(2), 215–222.
- Aydinalp, C., & Marinova, S. (2009). The effects of heavy metals on seed germination and plant growth on alfalfa plant (*Medicago sativa*). *Bulgarian Journal of Agricultural Science*, 15(4), 347–350.
- Bakırdere, S., Bölücek, C., & Yaman, M. (2016). Determination of contamination levels of Pb, Cd, Cu, Ni, and Mn caused by former lead mining gallery. *Environmental Monitoring and Assessment*, 188(3), 1–7.
- Bandara, T., Herath, I., Kumarathilaka, P., Hseu, Z.-Y., Ok, Y. S., & Vithanage, M. (2016). Efficacy of woody biomass and biochar for alleviating heavy metal bioavailability in serpentine soil. *Environmental Geochemistry and Health*. doi:10.1007/s10653-016-9842-0.
- Bandara, T., Herath, I., Kumarathilaka, P., Seneviratne, M., Seneviratne, G., Rajakaruna, N., et al. (2017). Role of woody biochar and fungal–bacterial co-inoculation on enzyme activity and metal immobilization in serpentine soil. *Journal of Soils and Sediments*, 17(3), 665–673. doi:10.1007/s11368-015-1243-y.
- Barceló, J., & Poschenrieder, C. (1990). Plant water relations as affected by heavy metal stress: A review. *Journal of Plant Nutrition*, 13(1), 1–37.
- Barceló, J., Poschenrieder, C., Andreu, I., & Gunse, B. (1986). Cadmium-induced decrease of water stress resistance in bush bean plants (*Phaseolus vulgaris* L. cv. Contender) I. Effects of Cd on water potential, relative water content, and cell wall elasticity. *Journal of Plant Physiology*, 125(1–2), 17–25.
- Bashmakov, D. I., Lukatkin, A. S., Revin, V. V., Duchovskis, P., Brazaitytė, A., & Baranauskis, K. (2005). Growth of maize seedlings affected by different concentrations of heavy metals. *Ekologija*, 3, 22–27.
- Basile, A., Sorbo, S., Conte, B., Cardi, M., & Esposito, S. (2013). Ultrastructural changes and heat shock proteins 70 induced by atmospheric pollution are similar to the effects observed under in vitro heavy metals stress in *Conocephalum conicum* (Marchantiales–Bryophyta). *Environmental Pollution*, 182, 209–216.
- Basta, N. T., Pantone, D., & Tabatabai, M. (1993). Path analysis of heavy metal adsorption by soil. *Agronomy Journal*, 85(5), 1054–1057.
- Baszyński, T. (2014). Interference of Cd²⁺ in functioning of the photosynthetic apparatus of higher plants. *Acta Societatis Botanicorum Poloniae*, 55(2), 291–304.
- Bénard, A., Rose, J., Hazemann, J.-L., Proux, O., Trotignon, L., Borschneck, D., et al. (2014). *Modelling the Pb release during Portland cement alteration*. arXiv:1402.2962.
- Bewley, J. D. (1997). Seed germination and dormancy. *The Plant Cell*, 9(7), 1055.
- Bolan, N. S., & Duraisamy, V. (2003). Role of inorganic and organic soil amendments on immobilisation and phytoavailability of heavy metals: A review involving specific case studies. *Soil Research*, 41(3), 533–555.
- Boyd, R. S., & Rajakaruna, N. (2013). Heavy metal tolerance. In D. Gibson (Ed.), *Oxford bibliographies in ecology*. New York: Oxford University Press.
- Branzini, A., & Zubillaga, M. S. (2012). Comparative use of soil organic and inorganic amendments in heavy metals stabilization. *Applied and Environmental Soil Science*, 2012, 7. doi:10.1155/2012/721032.
- Castaldi, P., Alberti, G., Merella, R., & Melis, P. (2005). Study of the organic matter evolution during municipal solid waste composting aimed at identifying suitable parameters for the evaluation of compost maturity. *Waste Management*, 25(2), 209–213. doi:10.1016/j.wasman.2004.12.011.
- Charest, C., & Ton Phan, C. (1990). Cold acclimation of wheat (*Triticum aestivum*): Properties of enzymes involved in proline metabolism. *Physiologia Plantarum*, 80(2), 159–168.
- Chen, C. T., Chen, L.-M., Lin, C. C., & Kao, C. H. (2001). Regulation of proline accumulation in detached rice leaves exposed to excess copper. *Plant Science*, 160(2), 283–290.
- Choudhary, N., Sairam, R., & Tyagi, A. (2005). Expression of Δ^1 -pyrroline-5-carboxylate synthetase gene during drought in rice (*Oryza sativa* L.). *Indian Journal of Biochemistry & Biophysics*, 42, 366–370.
- Chugh, L., & Sawhney, S. (1996). Effect of cadmium on germination, amylases and rate of respiration of germinating pea seeds. *Environmental Pollution*, 92(1), 1–5.
- Clemens, S. (2006). Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie*, 88(11), 1707–1719.
- Cokkizgin, A., & Cokkizgin, H. (2015). Effects of lead (PbCl₂) stress on germination of lentil (*Lens culinaris* Medic.) lines. *African Journal of Biotechnology*, 9(50), 8608–8612.
- Dago, Á., González, I., Ariño, C., Díaz-Cruz, J. M., & Esteban, M. (2014). Chemometrics applied to the analysis of

- induced phytochelatin in *Hordeum vulgare* plants stressed with various toxic non-essential metals and metalloids. *Talanta*, 118, 201–209.
- Davies, K. J. (2001). Degradation of oxidized proteins by the 20S proteasome. *Biochimie*, 83(3), 301–310.
- Davis, J. A. (1984). Complexation of trace metals by adsorbed natural organic matter. *Geochimica et Cosmochimica Acta*, 48(4), 679–691.
- De Lacerda, L. D., & Salomons, W. (2012). *Mercury from gold and silver mining: A chemical time bomb?*. Berlin: Springer Science & Business Media.
- Debeaujon, I., & Koornneef, M. (2000). Gibberellin requirement for Arabidopsis seed germination is determined both by testa characteristics and embryonic abscisic acid. *Plant Physiology*, 122(2), 415–424.
- Dermont, G., Bergeron, M., Mercier, G., & Richer-Lafleche, M. (2008). Soil washing for metal removal: A review of physical/chemical technologies and field applications. *Journal of Hazardous Materials*, 152(1), 1–31.
- Devi, R., Munjral, N., Gupta, A. K., & Kaur, N. (2013). Effect of exogenous lead on growth and carbon metabolism of pea (*Pisum sativum* L.) seedlings. *Physiology and Molecular Biology of Plants*, 19(1), 81–89.
- Diwan, H., Khan, I., Ahmad, A., & Iqbal, M. (2010). Induction of phytochelatin and antioxidant defence system in *Brassica juncea* and *Vigna radiata* in response to chromium treatments. *Plant Growth Regulation*, 61(1), 97–107.
- Downs, C. A., Coleman, J. S., & Heckathorn, S. A. (1999). The chloroplast 22-Ku heat-shock protein: A luminal protein that associates with the oxygen evolving complex and protects photosystem II during heat stress. *Journal of Plant Physiology*, 155(4), 477–487.
- Dua, A., & Sawhney, S. K. (1991). Effect of chromium on activities of hydrolytic enzymes in germinating pea seeds. *Environmental and Experimental Botany*, 31(2), 133–139.
- Dutta, P., Islam, M., & Mondal, S. (2014). Interactive effect of arsenic stress and seed phytate content on germination and seedling development of different vegetable crops. *Journal of Plant Physiology & Pathology*, 2(2), 2.
- Fabro, G., Kovács, I., Pavet, V., Szabados, L., & Alvarez, M. E. (2004). Proline accumulation and AtP5CS2 gene activation are induced by plant-pathogen incompatible interactions in Arabidopsis. *Molecular Plant-Microbe Interactions*, 17(4), 343–350.
- Falkowska, L., Reindl, A. R., Szumilo, E., Kwaśniak, J., Staniszevska, M., Bełdowska, M., et al. (2013). Mercury and chlorinated pesticides on the highest level of the food web as exemplified by herring from the Southern Baltic and African penguins from the zoo. *Water, Air, and Soil Pollution*, 224(5), 1–15.
- Fariduddin, Q., Khalil, R. R., Mir, B. A., Yusuf, M., & Ahmad, A. (2013). 24-Epibrassinolide regulates photosynthesis, antioxidant enzyme activities and proline content of *Cucumis sativus* under salt and/or copper stress. *Environmental Monitoring and Assessment*, 185(9), 7845–7856.
- Fu, J., Zhao, C., Luo, Y., Liu, C., Kyzas, G. Z., Luo, Y., et al. (2014). Heavy metals in surface sediments of the Jialu River, China: Their relations to environmental factors. *Journal of Hazardous Materials*, 270, 102–109.
- Fujii, H., Verslues, P. E., & Zhu, J.-K. (2007). Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in Arabidopsis. *The Plant Cell*, 19(2), 485–494.
- Fulton, D. C., Stettler, M., Mettler, T., Vaughan, C. K., Li, J., Francisco, P., et al. (2008). β -AMYLASE4, a noncatalytic protein required for starch breakdown, acts upstream of three active β -amylases in Arabidopsis chloroplasts. *The Plant Cell*, 20(4), 1040–1058.
- Gadepalle, V. P., Ouki, S. K., Herwijnen, R. V., & Hutchings, T. (2007). Immobilization of heavy metals in soil using natural and waste materials for vegetation establishment on contaminated sites. *Soil and Sediment Contamination: An International Journal*, 16(2), 233–251. doi:10.1080/15320380601169441.
- Gajewska, E., & Skłodowska, M. (2008). Differential biochemical responses of wheat shoots and roots to nickel stress: Antioxidative reactions and proline accumulation. *Plant Growth Regulation*, 54(2), 179–188.
- Gall, J. E., & Rajakaruna, N. (2013). The physiology, functional genomics, and applied ecology of heavy metal-tolerant Brassicaceae. In M. Lang (Ed.), *Brassicaceae: Characterization, functional genomics and health benefits* (pp. 121–148). New York: Nova.
- Gallego, S. M., Benavides, M. P., & Tomaro, M. L. (1996). Effect of heavy metal ion excess on sunflower leaves: Evidence for involvement of oxidative stress. *Plant Science*, 121(2), 151–159.
- Gianazza, E., Wait, R., Sozzi, A., Regondi, S., Saco, D., Labra, M., et al. (2007). Growth and protein profile changes in *Lepidium sativum* L. plantlets exposed to cadmium. *Environmental and Experimental Botany*, 59(2), 179–187.
- Gratão, P. L., Polle, A., Lea, P. J., & Azevedo, R. A. (2005). Making the life of heavy metal-stressed plants a little easier. *Functional Plant Biology*, 32(6), 481–494.
- Groot, S., & Karssen, C. (1987). Gibberellins regulate seed germination in tomato by endosperm weakening: A study with gibberellin-deficient mutants. *Planta*, 171(4), 525–531.
- Hadi, P., Gao, P., Barford, J. P., & McKay, G. (2013). Novel application of the nonmetallic fraction of the recycled printed circuit boards as a toxic heavy metal adsorbent. *Journal of Hazardous Materials*, 252, 166–170.
- Hall, J. (2002). Cellular mechanisms for heavy metal detoxification and tolerance. *Journal of Experimental Botany*, 53(366), 1–11.
- Han, D., Xiong, S., Tu, S., Liu, J., & Chen, C. (2015). Interactive effects of selenium and arsenic on growth, antioxidant system, arsenic and selenium species of *Nicotiana tabacum* L. *Environmental and Experimental Botany*, 117, 12–19.
- Hasanuzzaman, M., & Fujita, M. (2013). Exogenous sodium nitroprusside alleviates arsenic-induced oxidative stress in wheat (*Triticum aestivum* L.) seedlings by enhancing antioxidant defense and glyoxalase system. *Ecotoxicology*, 22(3), 584–596.
- He, J., Ren, Y., Chen, X., & Chen, H. (2014). Protective roles of nitric oxide on seed germination and seedling growth of rice (*Oryza sativa* L.) under cadmium stress. *Ecotoxicology and Environmental Safety*, 108, 114–119.
- Heckathorn, S. A., Mueller, J. K., LaGuidice, S., Zhu, B., Barrett, T., Blair, B., et al. (2004). Chloroplast small heat-shock proteins protect photosynthesis during heavy metal stress. *American Journal of Botany*, 91(9), 1312–1318.

- Herath, I., Iqbal, M. C. M., Al-Wabel, M. I., Abduljabbar, A., Ahmad, M., Usman, A. R. A., et al. (2017). Bioenergy-derived waste biochar for reducing mobility, bioavailability, and phytotoxicity of chromium in anthropized tannery soil. *Journal of Soils and Sediments*, 17(3), 731–740. doi:10.1007/s11368-015-1332-y.
- Herath, I., Kumarathilaka, P., Navaratne, A., Rajakaruna, N., & Vithanage, M. (2015). Immobilization and phytotoxicity reduction of heavy metals in serpentine soil using biochar. *Journal of Soils and Sediments*, 15(1), 126–138. doi:10.1007/s11368-014-0967-4.
- Hsu, Y., & Kao, C. (2003). Role of abscisic acid in cadmium tolerance of rice (*Oryza sativa* L.) seedlings. *Plant, Cell and Environment*, 26(6), 867–874.
- Hsu, Y. T., & Kao, C. H. (2005). Abscisic acid accumulation and cadmium tolerance in rice seedlings. *Physiologia Plantarum*, 124(1), 71–80.
- Hu, Y., Liu, X., Bai, J., Shih, K., Zeng, E. Y., & Cheng, H. (2013). Assessing heavy metal pollution in the surface soils of a region that had undergone three decades of intense industrialization and urbanization. *Environmental Science and Pollution Research*, 20(9), 6150–6159.
- Imran, M. A., Chaudhry, M., Khan, R., Ali, Z., & Mahmood, T. (2013). Toxicity of arsenic (As) on seed germination of sunflower (*Helianthus annuus* L.). *International Journal of Physical Sciences*, 8(17), 840–847.
- Inyang, M., Gao, B., Yao, Y., Xue, Y., Zimmerman, A. R., Pullammanappallil, P., et al. (2012). Removal of heavy metals from aqueous solution by biochars derived from anaerobically digested biomass. *Bioresource Technology*, 110, 50–56.
- Iqbal, N., Nazar, R., & Umar, S. (2016). Evaluating the importance of proline in cadmium tolerance and its interaction with phytohormones. In N. Iqbal, R. Nazar & N. A. Khan (Eds.), *Osmolytes and plants acclimation to changing environment: Emerging omics technologies* (pp. 129–153). Berlin: Springer.
- Iqbal, M. Z., Shafiq, M., & Athar, M. (2014). Phytotoxic effects of mercury on seed germination and seedling growth of *Albizia lebbek* (L.) Benth. (Leguminosae). *Advances in Environmental Research*, 3, 207–216.
- Irfan, M., Ahmad, A., & Hayat, S. (2014). Effect of cadmium on the growth and antioxidant enzymes in two varieties of *Brassica juncea*. *Saudi Journal of Biological Sciences*, 21(2), 125–131.
- Jackson, S. P., & Durocher, D. (2013). Regulation of DNA damage responses by ubiquitin and SUMO. *Molecular Cell*, 49(5), 795–807.
- Jiang, M., & Zhang, J. (2002). Water stress-induced abscisic acid accumulation triggers the increased generation of reactive oxygen species and up-regulates the activities of antioxidant enzymes in maize leaves. *Journal of Experimental Botany*, 53(379), 2401–2410.
- Kalai, T., Khamassi, K., Teixeira da Silva, J. A., Gouia, H., & Bettaieb Ben-Kaab, L. (2014). Cadmium and copper stress affect seedling growth and enzymatic activities in germinating barley seeds. *Archives of Agronomy and Soil Science*, 60(6), 765–783.
- Kang, J.-L., Zeng, Z.-J., & Liu, Y.-P. (2009). Effects of lead (Pb ~ (2+)) stress on seed germination and seedling growth of wheat. *Guangxi Agricultural Sciences*, 2, 009.
- Karmous, I., Bellani, L. M., Chaoui, A., El Ferjani, E., & Muccifora, S. (2015). Effects of copper on reserve mobilization in embryo of *Phaseolus vulgaris* L. *Environmental Science and Pollution Research*, 22(13), 10159–10165.
- Karmous, I., Chaoui, A., Jaouani, K., Sheehan, D., El Ferjani, E., Scoccianti, V., et al. (2014a). Role of the ubiquitin-proteasome pathway and some peptidases during seed germination and copper stress in bean cotyledons. *Plant Physiology and Biochemistry*, 76, 77–85.
- Karmous, I., Jaouani, K., El Ferjani, E., & Chaoui, A. (2014b). Responses of proteolytic enzymes in embryonic axes of germinating bean seeds under copper stress. *Biological Trace Element Research*, 160(1), 108–115.
- Karmous, I., Khadija, J., Chaoui, A., & El Ferjani, E. (2012). Proteolytic activities in *Phaseolus vulgaris* cotyledons under copper stress. *Physiology and Molecular Biology of Plants*, 18(4), 337–343.
- Kaul, S., Sharma, S., & Mehta, I. (2008). Free radical scavenging potential of L-proline: Evidence from in vitro assays. *Amino Acids*, 34(2), 315–320.
- Khan, A. (1968). Inhibition of gibberellic acid-induced germination by abscisic acid and reversal by cytokinins. *Plant Physiology*, 43(9), 1463.
- Khan, A., Kuek, C., Chaudhry, T., Khoo, C., & Hayes, W. (2000). Role of plants, mycorrhizae and phytochelators in heavy metal contaminated land remediation. *Chemosphere*, 41(1), 197–207.
- Kim, S., Lee, S., & Lee, I. (2012). Alteration of phytotoxicity and oxidant stress potential by metal oxide nanoparticles in *Cucumis sativus*. *Water, Air, and Soil Pollution*, 223(5), 2799–2806.
- Komatsu, S., Abbasi, F., Kobori, E., Fujisawa, Y., Kato, H., & Iwasaki, Y. (2005). Proteomic analysis of rice embryo: An approach for investigating G α protein-regulated proteins. *Proteomics*, 5(15), 3932–3941.
- Koomneef, M., Reuling, G., & Karssen, C. (1984). The isolation and characterization of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiologia Plantarum*, 61(3), 377–383.
- Kranner, I., & Colville, L. (2011). Metals and seeds: Biochemical and molecular implications and their significance for seed germination. *Environmental and Experimental Botany*, 72(1), 93–105.
- Kriz, A. L. (1989). Characterization of embryo globulins encoded by the maize Glb genes. *Biochemical Genetics*, 27(3–4), 239–251.
- Kumar, P. N., Dushenkov, V., Motto, H., & Raskin, I. (1995). Phytoextraction: The use of plants to remove heavy metals from soils. *Environmental Science and Technology*, 29(5), 1232–1238.
- Kumarathilaka, P., & Vithanage, M. (2017). Influence of *Gliciridia sepium* biochar on attenuate perchlorate-induced heavy metal release in serpentine soil. *Journal of Chemistry*, 2017, 8. doi:10.1155/2017/6180636.
- Küpper, H., Küpper, F., & Spiller, M. (1996). Environmental relevance of heavy metal-substituted chlorophylls using the example of water plants. *Journal of Experimental Botany*, 47(2), 259–266.
- Kuriakose, S. V., & Prasad, M. (2008). Cadmium stress affects seed germination and seedling growth in *Sorghum bicolor* (L.) Moench by changing the activities of hydrolyzing enzymes. *Plant Growth Regulation*, 54(2), 143–156.

- Labra, M., Gianazza, E., Waitt, R., Eberini, I., Sozzi, A., Regondi, S., et al. (2006). *Zea mays* L. protein changes in response to potassium dichromate treatments. *Chemosphere*, 62(8), 1234–1244.
- Lacerda, L. (1997). Global mercury emissions from gold and silver mining. *Water, Air, and Soil Pollution*, 97(3–4), 209–221.
- Li, C.-X., Feng, S.-L., Yun, S., Jiang, L.-N., Lu, X.-Y., & Hou, X.-L. (2007). Effects of arsenic on seed germination and physiological activities of wheat seedlings. *Journal of Environmental Sciences*, 19(6), 725–732.
- Lin, T.-Y., Wei, C.-C., Huang, C.-W., Chang, C.-H., Hsu, F.-L., & Liao, V. H.-C. (2016). Both phosphorus fertilizers and indigenous bacteria enhance arsenic release into groundwater in the arsenic-contaminated aquifers. *Journal of Agricultural and Food Chemistry*, 64(11), 2214–2222.
- Liu, J., Zhang, Y., Shi, P., & Chai, T. (2012). Effect of cadmium on seed germination and antioxidative enzymes activities in cotyledon of *Solanum nigrum* L. *Journal of Agro-Environment Science*, 31(5), 880–884.
- Mahdiah, S., Ghaderian, S. M., & Karimi, N. (2013). Effect of arsenic on germination, photosynthesis and growth parameters of two winter wheat varieties in Iran. *Journal of Plant Nutrition*, 36(4), 651–664.
- Maheshwari, R., & Dubey, R. S. (2008). Inhibition of ribonuclease and protease activities in germinating rice seeds exposed to nickel. *Acta Physiologiae Plantarum*, 30(6), 863–872.
- Maheshwari, R., & Dubey, R. (2009). Nickel-induced oxidative stress and the role of antioxidant defence in rice seedlings. *Plant Growth Regulation*, 59(1), 37–49.
- Mahmood, T., Islam, K., & Muhammad, S. (2007). Toxic effects of heavy metals on early growth and tolerance of cereal crops. *Pakistan Journal of Botany*, 39(2), 451.
- Malik, J. A., Goel, S., Kaur, N., Sharma, S., Singh, I., & Nayyar, H. (2012). Selenium antagonises the toxic effects of arsenic on mungbean (*Phaseolus aureus* Roxb.) plants by restricting its uptake and enhancing the antioxidative and detoxification mechanisms. *Environmental and Experimental Botany*, 77, 242–248.
- Mazhoudi, S., Chaoui, A., Ghorbal, M. H., & El Ferjani, E. (1997). Response of antioxidant enzymes to excess copper in tomato (*Lycopersicon esculentum*, Mill.). *Plant Science*, 127(2), 129–137.
- Mishra, S., & Dubey, R. S. (2005). Heavy metal toxicity induced alterations in photosynthetic metabolism in plants. In M. Pessaraki (Ed.), *Handbook of phytosynthesis*. Tucson: CRC Press.
- Mishra, S., & Dubey, R. S. (2006). Inhibition of ribonuclease and protease activities in arsenic exposed rice seedlings: Role of proline as enzyme protectant. *Journal of Plant Physiology*, 163(9), 927–936.
- Mishra, S., Jha, A., & Dubey, R. (2011). Arsenite treatment induces oxidative stress, upregulates antioxidant system, and causes phytochelatin synthesis in rice seedlings. *Protoplasma*, 248(3), 565–577.
- Mohamed, A. A., Castagna, A., Ranieri, A., & di Toppi, L. S. (2012). Cadmium tolerance in *Brassica juncea* roots and shoots is affected by antioxidant status and phytochelatin biosynthesis. *Plant Physiology and Biochemistry*, 57, 15–22.
- Muhammad, Z. I., Maria, K. S., Mohammad, A., Muhammad, S., Zia-Ur-Rehman, F., & Muhammad, K. (2015). Effect of mercury on seed germination and seedling growth of Mungbean (*Vigna radiata* (L.) Wilczek). *Journal of Applied Sciences and Environmental Management*, 19(2), 191–199.
- Nagajyoti, P. C., Lee, K. D., & Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: A review. *Environmental Chemistry Letters*, 8(3), 199–216. doi:10.1007/s10311-010-0297-8.
- Nambara, E., Akazawa, T., & McCourt, P. (1991). Effects of the gibberellin biosynthetic inhibitor uniconazol on mutants of Arabidopsis. *Plant Physiology*, 97(2), 736–738.
- Nasr, N. (2013). Germination and seedling growth of maize (*Zea mays* L.) seeds in toxicity of aluminum and nickel. *Merit Research Journal of Environmental Science and Toxicology*, 1(5), 110–113.
- Neilson, S., & Rajakaruna, N. (2015). Phytoremediation of agricultural soils: Using plants to clean metal-contaminated arable land. In A. A. Ansari, S. S. Gill, R. Gill, G. R. Lanza & L. Newman (Eds.), *Phytoremediation* (pp. 159–168). Berlin: Springer.
- Nicholson, F., Smith, S., Alloway, B., Carlton-Smith, C., & Chambers, B. (2003). An inventory of heavy metals inputs to agricultural soils in England and Wales. *Science of the Total Environment*, 311(1), 205–219.
- Østergaard, O., Finnie, C., Laugesen, S., Roepstorff, P., & Svensson, B. (2004). Proteome analysis of barley seeds: Identification of major proteins from two-dimensional gels (pI 4–7). *Proteomics*, 4(8), 2437–2447.
- Ouzounidou, G., Moustakas, M., & Eleftheriou, E. (1997). Physiological and ultrastructural effects of cadmium on wheat (*Triticum aestivum* L.) leaves. *Archives of Environmental Contamination and Toxicology*, 32(2), 154–160.
- Park, J. H., Lamb, D., Paneerselvam, P., Choppala, G., Bolan, N., & Chung, J.-W. (2011). Role of organic amendments on enhanced bioremediation of heavy metal(loids) contaminated soils. *Journal of Hazardous Materials*, 185(2–3), 549–574. doi:10.1016/j.jhazmat.2010.09.082.
- Patnaik, A., & Mohanty, B. (2013). Toxic effect of mercury and cadmium on germination and seedling growth of *Cajanus cajan* L. (pigeon pea). *Annals of Biological Research*, 4(3), 123–126.
- Perfus-Barbeoch, L., Leonhardt, N., Vavasseur, A., & Forestier, C. (2002). Heavy metal toxicity: Cadmium permeates through calcium channels and disturbs the plant water status. *The Plant Journal*, 32(4), 539–548.
- Prasad, M., & Strzałka, K. (1999). Impact of heavy metals on photosynthesis. In M. N. V. Prasad & J. Hagemeyer (Eds.), *Heavy metal stress in plants* (pp. 117–138). Berlin: Springer.
- Prodanovic, O., Prodanovic, R., Pristov, J. B., Mitrovic, A., & Radotic, K. (2016). Effect of cadmium stress on antioxidative enzymes during the germination of Serbian spruce [*Picea omorika* (Pan.) Purkyně]. *African Journal of Biotechnology*, 11(52), 11377–11385.
- Prost, R., & Yaron, B. (2001). Use of modified clays for controlling soil environmental quality. *Soil Science*, 166(12), 880–895.
- Rai, A., Tripathi, P., Dwivedi, S., Dubey, S., Shri, M., Kumar, S., et al. (2011). Arsenic tolerances in rice (*Oryza sativa*)

- have a predominant role in transcriptional regulation of a set of genes including sulphur assimilation pathway and antioxidant system. *Chemosphere*, 82(7), 986–995.
- Ranki, H., Sopanen, T., & Voutilainen, R. (1990). Localization of carboxypeptidase I in germinating barley grain. *Plant Physiology*, 93(4), 1449–1452.
- Rao, K. M., & Sresty, T. (2000). Antioxidative parameters in the seedlings of pigeonpea (*Cajanus cajan* (L.) Millspaugh) in response to Zn and Ni stresses. *Plant Science*, 157(1), 113–128.
- Rezaei, M., Sedghi, M., & Parmoon, G. (2013). Effect of HgCl₂ on the germination of hardened rapeseed. *Technical Journal of Engineering and Applied Sciences*, 3(13), 1162–1166.
- Rubio, M., Escrig, I., Martinez-Cortina, C., Lopez-Benet, F., & Sanz, A. (1994). Cadmium and nickel accumulation in rice plants. Effects on mineral nutrition and possible interactions of abscisic and gibberellic acids. *Plant Growth Regulation*, 14(2), 151–157.
- Salt, D. E., Prince, R. C., Pickering, I. J., & Raskin, I. (1995). Mechanisms of cadmium mobility and accumulation in Indian mustard. *Plant Physiology*, 109(4), 1427–1433.
- Sanal, F., Şeren, G., & Güner, U. (2014). Effects of arsenate and arsenite on germination and some physiological attributes of barley *Hordeum vulgare* L. *Bulletin of Environmental Contamination and Toxicology*, 92(4), 483–489.
- Sánchez-Virosta, P., Espín, S., García-Fernández, A., & Eeva, T. (2015). A review on exposure and effects of arsenic in passerine birds. *Science of the Total Environment*, 512, 506–525.
- Sankarganesh, K., Selvaraj, M., Baskaran, L., & Chidambaram, A. A. (2015). Chromium induced changes in Soybean (*Glycine max* L.) metabolism. *World Scientific News*, 16, 145–178.
- Saradhi, P. P. (1991). Proline accumulation under heavy metal stress. *Journal of Plant Physiology*, 138(5), 554–558.
- Saradhi, P. P., AliaArora, S., & Prasad, K. (1995). Proline accumulates in plants exposed to UV radiation and protects them against UV-induced peroxidation. *Biochemical and Biophysical Research Communications*, 209(1), 1–5.
- Saueia, C. H. R., Le Bourlegat, F. M., Mazzilli, B. P., & Fávoro, D. I. T. (2013). Availability of metals and radionuclides present in phosphogypsum and phosphate fertilizers used in Brazil. *Journal of Radioanalytical and Nuclear Chemistry*, 297(2), 189–195.
- Schat, H., Sharma, S. S., & Vooijs, R. (1997). Heavy metal-induced accumulation of free proline in a metal-tolerant and a nontolerant ecotype of *Silene vulgaris*. *Physiologia Plantarum*, 101(3), 477–482.
- Schützendübel, A., & Polle, A. (2002). Plant responses to abiotic stresses: Heavy metal-induced oxidative stress and protection by mycorrhization. *Journal of Experimental Botany*, 53(372), 1351–1365.
- Shah, K., & Dubey, K. (1995). Effect of cadmium on RNA level as well as activity and molecular forms of ribonuclease in growing rice seedlings. *Plant Physiology and Biochemistry*, 33(5), 577–584.
- Shah, K., & Dubey, R. (1998). Cadmium elevates level of protein, amino acids and alters activity of proteolytic enzymes in germinating rice seeds. *Acta Physiologiae Plantarum*, 20(2), 189–196.
- Shah, K., Kumar, R. G., Verma, S., & Dubey, R. (2001). Effect of cadmium on lipid peroxidation, superoxide anion generation and activities of antioxidant enzymes in growing rice seedlings. *Plant Science*, 161(6), 1135–1144.
- Shakeel, S. N., Haq, N. U., Heckathorn, S., & Luthe, D. (2012). Analysis of gene sequences indicates that quantity not quality of chloroplast small HSPs improves thermotolerance in C4 and CAM plants. *Plant Cell Reports*, 31(10), 1943–1957.
- Sharma, R. K., & Agrawal, M. (2005). Biological effects of heavy metals: An overview. *Journal of Environmental Biology*, 26(2), 301–313.
- Shen, G., Zhu, C., & Du, Q. Z. (2010). Genome-wide identification of PHYTOCHELATIN and PHYTOCH₂SYNTH domain-containing phytochelatin family from rice. *Electronic Journal of Biology*, 6(3), 73–79.
- Sheoran, I., Singal, H., & Singh, R. (1990). Effect of cadmium and nickel on photosynthesis and the enzymes of the photosynthetic carbon reduction cycle in pigeonpea (*Cajanus cajan* L.). *Photosynthesis Research*, 23(3), 345–351.
- Shri, M., Kumar, S., Chakrabarty, D., Trivedi, P. K., Mallick, S., Misra, P., et al. (2009). Effect of arsenic on growth, oxidative stress, and antioxidant system in rice seedlings. *Ecotoxicology and Environmental Safety*, 72(4), 1102–1110.
- Singh, H. P., Kaur, G., Batish, D. R., & Kohli, R. K. (2011). Lead (Pb)-inhibited radicle emergence in *Brassica campestris* involves alterations in starch-metabolizing enzymes. *Biological Trace Element Research*, 144(1–3), 1295–1301.
- Singh, D., Nath, K., & Sharma, Y. K. (2007). Response of wheat seed germination and seedling growth under copper stress. *Journal of Environmental Biology*, 28(2), 409.
- Singh, N., Singh, J., Kaur, L., Sodhi, N. S., & Gill, B. S. (2003). Morphological, thermal and rheological properties of starches from different botanical sources. *Food Chemistry*, 81(2), 219–231.
- Somashekaraiah, B., Padmaja, K., & Prasad, A. (1992). Phytotoxicity of cadmium ions on germinating seedlings of mung bean (*Phaseolus vulgaris*): Involvement of lipid peroxides in chlorophyll degradation. *Physiologia Plantarum*, 85(1), 85–89.
- Spurgeon, D., & Hopkin, S. (1996). Effects of variations of the organic matter content and pH of soils on the availability and toxicity of zinc to the earthworm *Eisenia fetida*. *Pedobiologia*, 40(1), 80–96.
- Srivastava, S., Akkarakaran, J. J., Suprasanna, P., & D'Souza, S. F. (2013). Response of adenine and pyridine metabolism during germination and early seedling growth under arsenic stress in *Brassica juncea*. *Acta Physiologiae Plantarum*, 35(4), 1081–1091.
- Srivastava, S., & Dubey, R. (2011). Manganese-excess induces oxidative stress, lowers the pool of antioxidants and elevates activities of key antioxidant enzymes in rice seedlings. *Plant Growth Regulation*, 64(1), 1–16.
- Srivastava, R. K., Pandey, P., Rajpoot, R., Rani, A., & Dubey, R. (2014). Cadmium and lead interactive effects on oxidative stress and antioxidative responses in rice seedlings. *Protoplasma*, 251(5), 1047–1065.
- Szabados, L., & Savouré, A. (2010). Proline: A multifunctional amino acid. *Trends in Plant Science*, 15(2), 89–97.

- Taiz, L., & Honigman, W. A. (1976). Production of cell wall hydrolyzing enzymes by barley aleurone layers in response to gibberellic acid. *Plant Physiology*, *58*(3), 380–386.
- Uchimiyama, M., Lima, I. M., Thomas Klasson, K., Chang, S., Wartelle, L. H., & Rodgers, J. E. (2010). Immobilization of heavy metal ions (CuII, CdII, NiII, and PbII) by broiler litter-derived biochars in water and soil. *Journal of Agricultural and Food Chemistry*, *58*(9), 5538–5544.
- Usman, A., Kuzyakov, Y., & Stahr, K. (2005). Effect of clay minerals on immobilization of heavy metals and microbial activity in a sewage sludge-contaminated soil (8 pp). *Journal of Soils and Sediments*, *5*(4), 245–252.
- Vázquez de Aldana, B., Gundel, P., García Criado, B., García Ciudad, A., García Sánchez, A., & Zabalgoitia, I. (2014). Germination response of endophytic *Festuca rubra* seeds in the presence of arsenic. *Grass & Forage Science*, *69*(3), 462–469.
- Waalkes, M. P. (2003). Cadmium carcinogenesis. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, *533*(1), 107–120.
- Wan, M., Wang, M., Zhou, F., & Yang, L. (2013). Effects of arsenic on seed germination of mung bean and black soybean. *Journal of Hubei University (Natural Science)*, *3*, 006.
- Wang, J., Zhao, F.-J., Meharg, A. A., Raab, A., Feldmann, J., & McGrath, S. P. (2002). Mechanisms of arsenic hyperaccumulation in *Pteris vittata*. Uptake kinetics, interactions with phosphate, and arsenic speciation. *Plant Physiology*, *130*(3), 1552–1561.
- Waters, E. R., Lee, G. J., & Vierling, E. (1996). Evolution, structure and function of the small heat shock proteins in plants. *Journal of Experimental Botany*, *47*(3), 325–338.
- Wu, G., Kang, H., Zhang, X., Shao, H., Chu, L., & Ruan, C. (2010). A critical review on the bio-removal of hazardous heavy metals from contaminated soils: Issues, progress, eco-environmental concerns and opportunities. *Journal of Hazardous Materials*, *174*(1), 1–8.
- Wuana, R. A., & Okieimen, F. E. (2011). Heavy metals in contaminated soils: A review of sources, chemistry, risks and best available strategies for remediation. *ISRN Ecology*. doi:10.5402/2011/402647.
- Xiang, L., Etxeberria, E., & den Ende, W. (2013). Vacuolar protein sorting mechanisms in plants. *FEBS Journal*, *280*(4), 979–993.
- Yan, D., Duermeyer, L., Leoveanu, C., & Nambara, E. (2014). The functions of the endosperm during seed germination. *Plant and Cell Physiology*, *55*(9), 1521–1533.
- Yan, H., Filardo, F., Hu, X., Zhao, X., & Fu, D. (2015). Cadmium stress alters the redox reaction and hormone balance in oilseed rape (*Brassica napus* L.) leaves. *Environmental Science and Pollution Research*, *23*(4), 1–12.
- Yang, S.-L., Lan, S.-S., & Gong, M. (2009). Hydrogen peroxide-induced proline and metabolic pathway of its accumulation in maize seedlings. *Journal of Plant Physiology*, *166*(15), 1694–1699.
- Yasur, J., & Rani, P. U. (2013). Environmental effects of nanosilver: Impact on castor seed germination, seedling growth, and plant physiology. *Environmental Science and Pollution Research*, *20*(12), 8636–8648.
- Yin, L., Colman, B. P., McGill, B. M., Wright, J. P., & Bernhardt, E. S. (2012). Effects of silver nanoparticle exposure on germination and early growth of eleven wetland plants. *PLoS ONE*, *7*(10), e47674.
- Yoshida, Y., Kiyosue, T., Katagiri, T., Ueda, H., Mizoguchi, T., Yamaguchi-Shinozaki, K., et al. (1995). Correlation between the induction of a gene for Δ^1 -pyrroline-5-carboxylate synthetase and the accumulation of proline in *Arabidopsis thaliana* under osmotic stress. *The Plant Journal*, *7*(5), 751–760.
- Zayneb, C., Bassem, K., Zeineb, K., Grubb, C. D., Noureddine, D., Hafedh, M., et al. (2015). Physiological responses of fenugreek seedlings and plants treated with cadmium. *Environmental Science and Pollution Research*, *22*(14), 10679–10689.
- Zeid, I. (2001). Responses of *Phaseolus vulgaris* chromium and cobalt treatments. *Biologia Plantarum*, *44*(1), 111–115.
- Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., et al. (2011). The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. *Environmental Pollution*, *159*(1), 84–91.
- Zhang, Q., & Riechers, D. E. (2004). Proteomic characterization of herbicide safener-induced proteins in the coleoptile of *Triticum tauschii* seedlings. *Proteomics*, *4*(7), 2058–2071.
- Zhang, S., Zhang, H., Qin, R., Jiang, W., & Liu, D. (2009). Cadmium induction of lipid peroxidation and effects on root tip cells and antioxidant enzyme activities in *Vicia faba* L. *Ecotoxicology*, *18*(7), 814–823.
- Zhi, Y., Deng, Z., Luo, M., Ding, W., Hu, Y., Deng, J., et al. (2015). Influence of heavy metals on seed germination and early seedling growth in *Eruca sativa* Mill. *American Journal of Plant Sciences*, *6*(05), 582.